

THE TROPICAL ANALOGUE OF POLAR CONES

STÉPHANE GAUBERT AND RICARDO D. KATZ

ABSTRACT. We study the max-plus or tropical analogue of the notion of polar: the polar of a cone represents the set of linear inequalities satisfied by its elements. We establish an analogue of the bipolar theorem, which characterizes all the inequalities satisfied by the elements of a tropical convex cone. We derive this characterization from a new separation theorem. We also establish variants of these results concerning systems of linear equalities.

1. INTRODUCTION

Max-plus or tropical algebra refers to the analogue of classical algebra obtained by considering the max-plus semiring $\mathbb{R}_{\max}$, which is the set $\mathbb{R} \cup \{-\infty\}$ equipped with the addition $a \oplus b := \max(a, b)$ and the multiplication $a \otimes b := a + b$.

Max-plus *convex cones* or *semimodules* are subsets V of $\mathbb{R}_{\max}^n$ stable by max-plus linear combinations, meaning that

$$(1) \quad \max(\lambda + x, \mu + y) \in V$$

for all $x, y \in V$ and $\lambda, \mu \in \mathbb{R}_{\max}$. Here, $\lambda + x$ denotes the vector with entries $\lambda + x_i$ and the “max” of two vectors is understood entrywise. Max-plus *convex sets* are subsets V of $\mathbb{R}_{\max}^n$ which satisfy (1) for all $x, y \in V$ and $\lambda, \mu \in \mathbb{R}_{\max}$ such that $\max(\lambda, \mu) = 0$.

Max-plus convex sets and cones have been studied by several authors with different motivations.

They were introduced by K. Zimmermann [Zim77] when studying discrete optimization problems.

Another motivation arose from the use by Maslov [Mas87] of a max-plus analogue of the superposition principle, which applies to the solutions of a stationary Hamilton-Jacobi equation whose Hamiltonian is convex in the adjoint variable. Hence, the space of solutions has a structure analogous to that of a linear space. This was one of the motivations for the development of the Idempotent Analysis [MS92, KM97] which includes the study, by Litvinov, Maslov, and Shpiz [LMS01] of the idempotent spaces, of which max-plus cones are special cases. See [LS07] for a recent development.

A third motivation comes from the algebraic approach to discrete event systems initiated by Cohen, Dubois, Quadrat and Viot [CDQV85]. The spaces that appear in the study of max-plus linear dynamical systems (reachable and observable spaces) are naturally equipped with structures of max-plus cones [CGQ99]. This motivated

2000 *Mathematics Subject Classification.* primary: 52A01, secondary: 16Y60, 46A55.

Key words and phrases. Max-plus semiring, max-plus convexity, tropical convexity, extremal convexity, $\mathbb{B}$ -convexity, duality, separation theorem, Farkas Lemma, semimodules, idempotent spaces.

The first author was partially supported by the joint CNRS-RFBR grant number 05-01-02807.

the study of max-plus cones or semimodules by Cohen, Quadrat, and the first author [CGQ01, CGQ04]. The results of [CGQ04] were completed in a work with Singer [CGQS05]. A development by the second author [Kat07] of the work on discrete event systems includes a max-plus cone based synthesis of controllers. See also [GK04].

The theory of max-plus convex cones, or tropical convex sets, has recently been developed in relation to tropical geometry. The tropical analogues of polytopes were considered by Develin and Sturmfels [DS04], who established a remarkable connection with phylogenetic analysis, showing that tree metrics may be thought of in terms of tropical polytopes. Tropical convexity was further studied by Joswig [Jos05] and Develin and Yu [DY07]. See [JSY07] for a new development.

Another interest in the max-plus analogues of convex cones comes from abstract convex analysis [Sin97], as can be seen in the work of Nitica and Singer [NS07a, NS07b, NS07c]. Independently, Briec, Horvath and Rubinov [BH04, BHR05] introduced and studied the notion of $\mathbb{B}$ -convexity which is another name for max-plus convexity. See also [AR06].

These motivations led to the proof of max-plus analogues of classical results such as: Hahn-Banach theorem either in analytic (extension of linear forms) or geometric (separation) form [Zim77, SS92, CGQ01, DS04, CGQ04, BHR05, CGQS05], Minkowski theorem (generation of a closed convex set by its extreme points) [Hel88, GK06, BSS07, GK07], and Helly and Carathéodory type theorems [BH04, GS07, GM08]. However, some duality results are still missing, because the idempotency of addition creates difficulties which are absent in the classical theory. For example, in the classical theory there exists a bijective correspondence between the facets of a convex cone and the extreme rays of its polar cone. This kind of properties, relating internal and external representations of convex cones, are not yet well understood in the max-plus case.

Indeed, classical duality theory relates a convex cone with the set of linear inequalities that its points satisfy. In the max-plus setting, the *polar* of a max-plus convex cone $V \subset \mathbb{R}_{\max}^n$ can be defined as

$$V^\circ = \{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid \max_i (f_i + x_i) \leq \max_j (g_j + x_j), \forall x \in V\}.$$

Conversely, we may consider the set of points satisfying a given set of inequalities. This leads us to define, for all $W \subset (\mathbb{R}_{\max}^n)^2$, a “dual” polar cone:

$$W^\diamond = \{x \in \mathbb{R}_{\max}^n \mid \max_i (f_i + x_i) \leq \max_j (g_j + x_j), \forall (f, g) \in W\}.$$

It follows from the separation theorem of [Zim77, SS92, CGQS05] that a closed max-plus convex cone V is characterized by its polar cone:

$$(V^\circ)^\diamond = V.$$

It is natural to ask which subsets W of $(\mathbb{R}_{\max}^n)^2$ arise as polars of cones, or equivalently, which subsets are of the form $W = (U^\circ)^\diamond$ for some $U \subset (\mathbb{R}_{\max}^n)^2$. The main result of this paper is the following analogue of the bipolar theorem:

Theorem 1 (Bipolar). *A subset $W \subset (\mathbb{R}_{\max}^n)^2$ is the polar of a cone if, and only if, the following conditions are satisfied:*

- (i) W is a closed max-plus convex cone,
- (ii) $f \leq g \implies (f, g) \in W$,
- (iii) $(f, g) \in W, (g, h) \in W \implies (f, h) \in W$.

This follows from Theorem 10 below. We deduce this result from a new separation theorem, Theorem 8, which is in some sense a dual of the separation theorem of [CGQS05].

We shall first establish similar results when the scalars form a complete semiring which is reflexive in the sense of [CGQ04]. This applies in particular to the completion of the max-plus semiring $\overline{\mathbb{R}}_{\max}$. Thanks to completeness and reflexivity properties, we shall obtain the separation theorem by algebraic and order theoretical arguments, relying on residuation theory (Galois connection in lattices). Then, we shall derive the results in the case of the max-plus semiring by a perturbation argument.

The notion of polar is illustrated in Figures 1 and 2 below. Figure 1 represents a max-plus convex cone V and eight max-plus linear inequalities composing a set W such that $V = W^\circ$. For visibility reasons, we have depicted completely only one of these inequalities, which is also shown separately. Detailed explanations of the constructions of Figure 1 can be found in Example 3.

We shall show that if W is a finite subset of $(\mathbb{R}_{\max}^n)^2$, representing a finite set of linear inequalities, then there are in general elements in the transitive closure of W which are not linear combinations of W . This means that there are some inequalities which can be logically deduced from the finite set of inequalities

$$\max_i (f_i + x_i) \leq \max_j (g_j + x_j) , \quad (f, g) \in W$$

but which cannot be obtained by a linear combination of these inequalities. Hence, Theorem 1 shows that there is no direct analogue of Farkas lemma.

This is related to the difficulties in defining the faces of tropical polyhedra, due to the obstructions mentioned in [GK07] and [DY07].

Some properties concerning max-plus convex cones are analogous to the case of cones in classical convexity. However, max-plus convex cones also have some features which are reminiscent of classical linear spaces. This is because in max-plus algebra, an inequality $\max_i (f_i + x_i) \leq \max_j (g_j + x_j)$ can always be rewritten as an equality, $\max_i (\max(f_i, g_i) + x_i) = \max_j (g_j + x_j)$. This point of view leads to define [GP97] the orthogonal of a max-plus convex cone V ,

$$V^\perp = \{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid \max_i (f_i + x_i) = \max_j (g_j + x_j), \forall x \in V\} .$$

Such an orthogonal is a congruence of semimodules, in the sense that will be defined in Section 3. Like in the case of linear inequalities, we may also consider the set of points satisfying a given set of linear equalities: for $W \subset (\mathbb{R}_{\max}^n)^2$ define

$$W^\top = \{x \in \mathbb{R}_{\max}^n \mid \max_i (f_i + x_i) = \max_j (g_j + x_j), \forall (f, g) \in W\} .$$

As a consequence of the separation theorem of [Zim77, SS92, CGQS05] we have

$$(V^\perp)^\top = V ,$$

if V is a closed cone, a result which was already stated in [GP97] in the particular case of finitely generated cones. We shall prove, as a variant of our main theorem, the following biorthogonal theorem. Like the bipolar theorem above (Theorem 1), it also follows from Theorem 10.

Theorem 2 (Biorthogonal). *A subset $W \subset (\mathbb{R}_{\max}^n)^2$ is the orthogonal of a cone if, and only if, the following conditions are satisfied:*

- (i) *W is a closed max-plus convex cone,*

- (ii) $(f, f) \in W$, $\forall f \in \mathbb{R}_{\max}^n$,
- (iii) $(f, g) \in W \implies (g, f) \in W$,
- (iv) $(f, g) \in W$, $(g, h) \in W \implies (f, h) \in W$.

Congruences are used in [LGKL08] to solve some observability problems for max-plus linear discrete event systems, i.e. to reconstruct some information on the state of the system from observations. The present biorthogonality theorem is used there to construct dynamic observers: in fact, cones are simpler to manipulate than congruences, due to the absence of “doubling” of the dimension, duality allows us to reduce algorithmic problems concerning congruences to algorithmic problems concerning cones. The duality results of the present paper may also be useful in the recent application of max-plus polyhedra to static analysis by abstract interpretation [AGG08].

2. PRELIMINARIES

In this section we recall some notions and results concerning semimodules over idempotent semirings which we shall need throughout this paper. We refer the reader to [BCOQ92, CGQ04, BJ72] for more background.

A monotone map $F : T \rightarrow S$ between two ordered sets $(T, \leq)$ and $(S, \leq)$ is said to be *residuated* if for each $s \in S$ the least upper bound of the set $\{t \in T \mid F(t) \leq s\}$ exists and belongs to it. When F is residuated, the map $F^\sharp : S \rightarrow T$ defined by $F^\sharp(s) := \vee \{t \in T \mid F(t) \leq s\}$, where $\vee Q$ denotes the least upper bound of Q , is called the *residual* of F . We say that F is *continuous* if $F(\vee Q) = \vee F(Q)$ for all $Q \subset T$, where $F(Q) := \{F(t) \mid t \in Q\}$. The term continuous refers to the Scott topology [GHK⁺03], which is a (non-Hausdorff) topology well adapted to the study of completed ordered algebraic structures. When T and S are complete ordered sets, meaning that any of their subsets admits a least upper bound, it can be shown that $F : T \rightarrow S$ is residuated if, and only if, it is continuous (see [BJ72, Th. 5.2] or [BCOQ92, Th. 4.50]).

Recall that any idempotent semiring $(\mathcal{K}, \oplus, \otimes, \varepsilon, e)$, where ε and e represent the neutral elements for addition $\oplus$ and multiplication $\otimes$ respectively, can be equipped with the *natural* order relation: $a \leq b \iff a \oplus b = b$, for which $\vee \{a, b\} = a \oplus b$. We say that an idempotent semiring $\mathcal{K}$ is *complete* if it is complete as a naturally ordered set, and if the left and right multiplications, $L_a : \mathcal{K} \rightarrow \mathcal{K}$, $L_a(b) = a \otimes b$ and $R_a : \mathcal{K} \rightarrow \mathcal{K}$, $R_a(b) = b \otimes a$, are continuous.

Given a semiring $(\mathcal{K}, \oplus, \otimes, \varepsilon_{\mathcal{K}}, e)$, a right $\mathcal{K}$ -*semimodule* is a commutative monoid $(X, \hat{\oplus})$, with neutral element ε_X , equipped with a map $X \times \mathcal{K} \rightarrow X$, $(x, \lambda) \rightarrow x \cdot \lambda$, (right action), which satisfies:

$$\begin{aligned} (x \hat{\oplus} z) \cdot \lambda &= x \cdot \lambda \hat{\oplus} z \cdot \lambda, & x \cdot (\lambda \oplus \mu) &= x \cdot \lambda \hat{\oplus} x \cdot \mu, & x \cdot (\lambda \otimes \mu) &= (x \cdot \lambda) \cdot \mu, \\ x \cdot \varepsilon_{\mathcal{K}} &= \varepsilon_X, & \varepsilon_X \cdot \lambda &= \varepsilon_X, & x \cdot e &= x, \end{aligned}$$

for all $x, z \in X$ and $\lambda, \mu \in \mathcal{K}$. Henceforth, the semimodule addition will be denoted by $\oplus$ (instead of $\hat{\oplus}$), like the semiring addition, and we will use concatenation to denote both the multiplication of $\mathcal{K}$ and the right action. Also for simplicity, we will denote by ε both the zero element $\varepsilon_{\mathcal{K}}$ of $\mathcal{K}$ and the zero element ε_X of X . Left $\mathcal{K}$ -semimodules are defined dually. A *subsemimodule* of X is a subset $V \subset X$ such that $x\lambda \oplus z\mu \in V$ for all $x, z \in V$ and $\lambda, \mu \in \mathcal{K}$. Note that a right (or left) $\mathcal{K}$ -semimodule $(X, \oplus)$ is necessarily idempotent when $\mathcal{K}$ is idempotent. In this case, X is said to be *complete*, if it is complete as a naturally ordered set, and if for all

$u \in X$ and $\lambda \in \mathcal{K}$, the maps $R_\lambda : X \rightarrow X$, $z \mapsto z\lambda$ and $L_u : \mathcal{K} \rightarrow X$, $\mu \mapsto u\mu$, are both continuous. Then, we can define

$$u \backslash x := L_u^\#(x) = \vee \{\mu \in \mathcal{K} \mid u\mu \leq x\} \quad \text{and} \quad x/\lambda := R_\lambda^\#(x) = \vee \{z \in X \mid z\lambda \leq x\},$$

for all $x, u \in X$ and $\lambda \in \mathcal{K}$. Note that by definition, we have $u(u \backslash x) \leq x$ and $(x/\lambda)\lambda \leq x$. A subsemimodule V of X is complete, if $\vee Q \in V$ for all $Q \subset V$. It is convenient to recall the following universal separation theorem for complete semimodules.

Theorem 3 ([CGQ04, Th. 8]). *Let $V \subset X$ be a complete subsemimodule. Assume that $x \notin V$. Then,*

$$v \in V \implies v \backslash P_V(x) = v \backslash x$$

and

$$x \backslash P_V(x) \neq x \backslash x,$$

where $P_V(x) := \vee \{v \in V \mid v \leq x\}$.

Example 1. The set $\mathbb{R}_{\max}^n$ is a right $\mathbb{R}_{\max}$ -semimodule when it is equipped with the usual operations: $(x \oplus z)_i := x_i \oplus z_i$ and $(x\lambda)_i := x_i\lambda$ for all $x, z \in \mathbb{R}_{\max}^n$ and $\lambda \in \mathbb{R}_{\max}$. Since $\mathbb{R}_{\max}$ is not a complete semiring, it will be convenient to consider the *completed* max-plus semiring $\overline{\mathbb{R}}_{\max}$, which is obtained by adjoining the element $+\infty$ to $\mathbb{R}_{\max}$.

In order to state separation theorems with the usual scalar product form, we next recall the notion of dual pair of [CGQ04], which is analogous to the one that arises in the theory of topological vector spaces (see for example [Bou64]). As usual, a map between right (resp. left) $\mathcal{K}$ -semimodules is right (resp. left) *linear* if it preserves finite sums and commutes with the action of scalars. We call *pre-dual pair* for a complete idempotent semiring $\mathcal{K}$, a complete right $\mathcal{K}$ -semimodule X together with a complete left $\mathcal{K}$ -semimodule Y equipped with a bracket $\langle \cdot \mid \cdot \rangle$ from $Y \times X$ to $\mathcal{K}$, such that, for all $x \in X$ and $y \in Y$, the maps $y \mapsto \langle y \mid x \rangle$ and $x \mapsto \langle y \mid x \rangle$ are respectively left and right linear, and continuous. We shall denote by (Y, X) this pre-dual pair. We say that Y *separates* X if

$$(\forall y \in Y, \langle y \mid x \rangle = \langle y \mid x' \rangle) \implies x = x',$$

and that X *separates* Y if

$$(\forall x \in X, \langle y \mid x \rangle = \langle y' \mid x \rangle) \implies y = y'.$$

A pre-dual pair (Y, X) such that X separates Y and Y separates X is called *dual pair*.

When X is a complete right $\mathcal{K}$ -semimodule, we call *opposite* semimodule of X the left $\mathcal{K}$ -semimodule X^{op} whose set of elements is X equipped with the addition $(x, u) \mapsto \wedge \{x, u\}$ and the left action $(\lambda, x) \mapsto x/\lambda$. Here, $\wedge \{x, u\}$ is the greatest lower bound of $\{x, u\}$ for the natural order of X . The fact that X^{op} is a complete left $\mathcal{K}$ -semimodule follows from properties of the residual map x/λ (see [CGQ04] for details). In particular, the complete idempotent semiring $\mathcal{K}$ can be thought of as a right $\mathcal{K}$ -semimodule, and so the same construction and notation apply to $\mathcal{K}$.

Theorem 4 ([CGQ04, Th. 22]). *Let X be a complete right $\mathcal{K}$ -semimodule. Then, the semimodules X^{op} and X form a dual pair for the bracket $X^{\text{op}} \times X \rightarrow \mathcal{K}^{\text{op}}$, $(y, x) \mapsto \langle y \mid x \rangle = x \backslash y$.*

Given a pre-dual pair (Y, X) for $\mathcal{K}$ and an arbitrary element $\varphi \in \mathcal{K}$, consider the maps:

$$\begin{aligned} X &\rightarrow Y, \quad x \mapsto \neg x = \vee\{y \in Y \mid \langle y \mid x \rangle \leq \varphi\}, \\ Y &\rightarrow X, \quad y \mapsto y^- = \vee\{x \in X \mid \langle y \mid x \rangle \leq \varphi\}. \end{aligned}$$

Note that if the dual pair for $\mathcal{K}^{\text{op}}$ of Theorem 4 is used, then the suprema in the definition above are actually infima (in terms of the order of $\mathcal{K}$) and the inequalities are reversed. The elements of X and Y of the form y^- and $\neg x$, respectively, are called *closed*. For $\lambda \in \mathcal{K}$, we define

$$\lambda^- := \lambda \setminus \varphi = \vee\{\mu \in \mathcal{K} \mid \lambda\mu \leq \varphi\} \quad \text{and} \quad \neg\lambda := \varphi/\lambda = \vee\{\mu \in \mathcal{K} \mid \mu\lambda \leq \varphi\}.$$

A complete idempotent semiring $\mathcal{K}$ is *reflexive* if there exists $\varphi \in \mathcal{K}$ such that $\neg(\lambda^-) = \lambda$ and $(\neg\lambda)^- = \lambda$ for all $\lambda \in \mathcal{K}$. We shall need the following proposition.

Proposition 1 ([CGQ04, Prop. 32 and Prop. 27]). *If $\mathcal{K}$ is reflexive and (Y, X) is a pre-dual pair for which Y separates X , then, all the elements of X are closed. Moreover, for all $x, u \in X$ it is satisfied:*

$$(2) \quad u \setminus x = \langle \neg x \mid u \rangle^-.$$

Example 2. The completed max-plus semiring $\overline{\mathbb{R}}_{\max}$ is reflexive. If we take $\varphi = 0$, then $\neg\lambda = \lambda^- = -\lambda$ for all $\lambda \in \overline{\mathbb{R}}_{\max}$. If $\overline{\mathbb{R}}_{\max}^n$ is equipped with the bracket $\langle y \mid x \rangle := \oplus_i y_i x_i$, then (2) becomes $u \setminus x = -(\oplus_i u_i(-x_i))$ for all $u, x \in \overline{\mathbb{R}}_{\max}^n$.

The results above are related to the max-plus analogues of the representation theorem for linear forms and the analytic form of the Hahn-Banach theorem of [CGQ04] which extend the corresponding results of [LMS01]. For a detailed presentation and examples of pre-dual pairs and reflexive semirings, we refer the reader to [CGQ04].

3. SEPARATION THEOREMS

Given a right $\mathcal{K}$ -semimodule X , a subset W of X^2 is called a *pre-congruence* if it is a subsemimodule such that

$$(f, f) \in W, \quad \forall f \in X$$

and

$$(f, g) \in W, \quad (g, h) \in W \implies (f, h) \in W.$$

If in addition a pre-congruence W verifies

$$(f, g) \in W \implies (g, f) \in W,$$

then W is said to be a *congruence*. In other words, a pre-congruence (resp. congruence) is a pre-order (resp. equivalence) relation on X which has a semimodule structure when it is thought of as a subset of the semimodule X^2 . A pre-congruence W which satisfies the property

$$(3) \quad f \leq g \implies (f, g) \in W,$$

is called a *polar cone*.

Remark 1. When W is a pre-congruence, note that (3) is equivalent to

$$(4) \quad f \leq g, \quad (g, h) \in W \implies (f, h) \in W.$$

Indeed, clearly (4) implies (3) because $(g, g) \in W$ for all $g \in X$. Conversely, since W is a pre-congruence, it follows that $(f, g) \in W$ and $(g, h) \in W$ imply $(f, h) \in W$, and so (3) implies (4).

The definition above is motivated by the classical notion of polar cones. Recall that if $V \subset \mathbb{R}^n$ is a (classical) convex cone, then its polar is defined as

$$\{f \in \mathbb{R}^n \mid \sum_i f_i x_i \leq 0, \forall x \in V\},$$

where $f_i x_i$ denotes the usual multiplication of the real numbers f_i and x_i . The direct extension of this definition to $X = \mathbb{R}_{\max}^n$, i.e.

$$\{f \in \mathbb{R}_{\max}^n \mid \oplus_i f_i x_i \leq \varepsilon, \forall x \in V\},$$

where $f_i x_i$ denotes $f_i \otimes x_i = f_i + x_i$, the usual addition of real numbers, is not convenient because this set is usually trivial independently of $V \subset \mathbb{R}_{\max}^n$. However, as it was mentioned in the introduction, the notion of polar can be extended to the max-plus algebra framework if we consider pairs of vectors instead of individual vectors, i.e. if we define the polar of V by

$$(5) \quad W := \{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid \oplus_i f_i x_i \leq \oplus_j g_j x_j, \forall x \in V\}.$$

Observe that any set of this form is a polar cone of X^2 in the sense defined above. However, not all polar cones have this form (i.e. are the polar of a cone V), take for example $W = \{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid g = \varepsilon \implies f = \varepsilon\}$.

Example 3. Let $\mathcal{K} = \mathbb{R}_{\max}$ and $X = \mathbb{R}_{\max}^3$. Consider the semimodule $V \subset X$ generated by the vectors $a = (1, 0, 2)^T$, $b = (1, 1, 0)^T$ and $c = (2, 4, 0)^T$, i.e. the semimodule composed of all the max-plus linear combinations of these three vectors. This semimodule is represented by the bounded dark gray region of Figure 1 together with the segments joining the points a and c to it. In this figure, every element of V with finite entries is projected orthogonally onto any plane orthogonal to the main diagonal $(1, 1, 1)^T$ of $\mathbb{R}^3$. Note that vectors that are proportional in the max-plus sense are sent to the same point, so this gives a faithful image of V .

If we define the semimodule $W \subset X^2$ by (5), then it can be checked that W is a polar cone of X^2 . In other words, this polar cone is composed of all the pairs of vectors (f, g) such that the corresponding max-plus half-space $\{x \in \mathbb{R}_{\max}^3 \mid \oplus_i f_i x_i \leq \oplus_j g_j x_j\}$ contains V . For example, if we take $f = (2, \varepsilon, \varepsilon)^T$ and $g = (\varepsilon, 0, 3)^T$, then $(f, g) \in W$ because a, b, c and hence V are contained in the half-space $\{x \in \mathbb{R}_{\max}^3 \mid 2x_1 \leq x_2 \oplus 3x_3\}$, which is represented by the unbounded light gray region of Figure 1.

In this case there exist eight max-plus linear inequalities $\oplus_i f_i x_i \leq \oplus_j g_j x_j$ which satisfy that V is the intersection of the corresponding half-spaces. These inequalities, which are associated with elements of W as explained above, are the following: $x_2 \leq 2x_1$, $x_2 \leq 4x_3$, $x_3 \leq 2x_2$, $x_3 \leq 1x_1$, $x_1 \leq 1x_2$, $x_1 \leq 2x_3$, $1x_1 \leq 1x_2 \oplus x_3$ and $2x_1 \leq x_2 \oplus 3x_3$. The boundaries of the corresponding eight half-spaces have been represented in Figure 1. Only the half-space corresponding to the last inequality has been depicted completely (note that this inequality is the one associated with the pair of vectors $(f, g) \in W$ considered above).

Since the purpose of this section is to establish separation theorems for complete congruences and polar cones, in the rest of this section we will assume that $\mathcal{K}$ is a complete idempotent semiring and that X is a complete right $\mathcal{K}$ -semimodule.




FIGURE 1. The semimodule V , a supporting half-space, and the polar of V .

Given a complete pre-congruence $W \subset X^2$, for $g \in X$, we set

$$(6) \quad \hat{g} := \vee \{f \in X \mid (f, g) \in W\}.$$

Observe that $(\hat{g}, g) \in W$ because W is complete. Moreover, since $(g, g) \in W$, it follows that $g \leq \hat{g}$ for all $g \in X$.

When W is a congruence, $\hat{g}$ is nothing but the maximal element in the equivalence class of g modulo W . Therefore, in this case we have the following obvious properties, which hold for all $g, f \in X$,

$$(7) \quad g \leq \hat{g} = \hat{\hat{g}},$$

$$(8) \quad (f, g) \in W \iff \hat{f} = \hat{g}.$$

We shall use the fact that

$$(9) \quad f \leq g \implies \hat{f} \leq \hat{g}.$$

Indeed, if $f \leq g$ then $g = f \oplus g$, and if $h \in X$ is such that $(h, f) \in W$, we have $(h \oplus g, g) = (h \oplus g, f \oplus g) = (h, f) \oplus (g, g) \in W$ because W is a pre-congruence. It follows that $\hat{g} \geq h \oplus g \geq h$. Since this holds for all $h \in X$ such that $(h, f) \in W$, we deduce that $\hat{g} \geq \hat{f}$, which shows (9).

Lemma 1. *Let $W \subset X^2$ be a complete pre-congruence. Then,*

$$(f, g) \in W \implies g \setminus \hat{h} \leq f \setminus \hat{h}$$

for all $h \in X$.

Proof. Let $\lambda := g \setminus \hat{h}$, so that $\hat{h} \geq g\lambda$, or equivalently $\hat{h} = \hat{h} \oplus g\lambda$. Since W is a pre-congruence, from $(\hat{h} \oplus f\lambda, \hat{h}) = (\hat{h} \oplus f\lambda, \hat{h} \oplus g\lambda) = (\hat{h}, \hat{h}) \oplus (f, g)\lambda \in W$ and $(\hat{h}, h) \in W$, we deduce that $(\hat{h} \oplus f\lambda, h) \in W$ and so $f\lambda \leq \hat{h} \oplus f\lambda \leq \hat{h}$. Hence, $f \setminus \hat{h} \geq \lambda$, i.e. $f \setminus \hat{h} \geq g \setminus \hat{h}$. $\square$

Now it is possible to prove the following separation theorem for complete polar cones.

Theorem 5. *Let $W \subset X^2$ be a complete polar cone. Assume that $s, t \in X$ are such that $(s, t) \notin W$. Then, there exists $x \in X$ such that*

$$(f, g) \in W \implies g \setminus x \leq f \setminus x$$

and

$$t \setminus x \not\leq s \setminus x.$$

Remark 2. In other words, Theorem 5 says that the set

$$M := \{(f, g) \in X^2 \mid g \setminus x \leq f \setminus x\}$$

separates the complete polar cone W and the point (s, t) : $W \subset M$ and $(s, t) \notin M$.

Proof of Theorem 5. We will show that the assertion of the theorem holds with $x := \hat{t}$.

In the first place, note that if $x = \hat{t}$, then by Lemma 1 the first assertion of the theorem is satisfied. Assume that $t \setminus \hat{t} \leq s \setminus \hat{t}$. Then, we get $s(t \setminus \hat{t}) \leq s(s \setminus \hat{t}) \leq \hat{t}$. Since $\hat{t} \geq t$, we have $t \setminus \hat{t} \geq e$, and so $s \leq \hat{t}$. Thus, from $s \leq \hat{t}$ and $(\hat{t}, t) \in W$, it follows that $(s, t) \in W$ because W is a polar cone, which contradicts our assumption. $\square$

In the case of complete congruences, we have the following separation theorem.

Theorem 6. *Let $W \subset X^2$ be a complete congruence. Assume that $s, t \in X$ are such that $(s, t) \notin W$. Then, there exists $x \in X$ such that*

$$(f, g) \in W \implies f \setminus x = g \setminus x$$

and

$$s \setminus x \neq t \setminus x.$$

Proof. Since $(s, t) \notin W$, by (8) we have $\hat{s} \neq \hat{t}$, hence, $\hat{s} \not\leq \hat{t}$ and/or $\hat{t} \not\leq \hat{s}$. Assume for instance that $\hat{t} \not\leq \hat{s}$. Then, we claim that

$$s \setminus \hat{s} \neq t \setminus \hat{s}.$$

Indeed, if $s \setminus \hat{s} = t \setminus \hat{s}$, we get $t(s \setminus \hat{s}) = t(t \setminus \hat{s}) \leq \hat{s}$. Since $\hat{s} \geq s$, we have $s \setminus \hat{s} \geq e$, and so $t \leq \hat{s}$. From (7) and (9), it follows that $\hat{t} \leq \hat{s} = \hat{s}$, which contradicts our assumption, so this proves our claim. Finally, as $(f, g) \in W$ implies that $(g, f) \in W$ because W is a congruence, by Lemma 1 the assertion of the theorem holds with $x := \hat{s}$. $\square$

To recover separation theorems with the usual scalar product form, we need to consider reflexive semirings.

Corollary 1. *Let $W \subset X^2$ be a complete congruence and (Y, X) be a pre-dual pair for a reflexive semiring $\mathcal{K}$ such that Y separates X . If $(s, t) \notin W$, then, there exists $y \in Y$ such that*

$$(f, g) \in W \implies \langle y \mid f \rangle = \langle y \mid g \rangle$$

and

$$\langle y \mid s \rangle \neq \langle y \mid t \rangle .$$

Proof. By Theorem 6 and Proposition 1 it follows that

$$(f, g) \in W \implies \langle \neg x \mid f \rangle^- = \langle \neg x \mid g \rangle^-$$

and

$$\langle \neg x \mid s \rangle^- \neq \langle \neg x \mid t \rangle^- ,$$

for some $x \in X$. Since $\mathcal{K}$ is reflexive, the map $\lambda \mapsto \lambda^-$ is injective, and thus we have

$$(f, g) \in W \implies \langle \neg x \mid f \rangle = \langle \neg x \mid g \rangle$$

and

$$\langle \neg x \mid s \rangle \neq \langle \neg x \mid t \rangle .$$

Therefore, the assertion of the theorem holds with $y := \neg x$. $\square$

The following corollary of Theorem 5 and Proposition 1 can be proved along similar lines, using the fact that

$$\lambda_1 \leq \lambda_2 \implies \neg \lambda_2 \leq \neg \lambda_1 , \lambda_2^- \leq \lambda_1^-$$

for all $\lambda_1, \lambda_2 \in \mathcal{K}$.

Corollary 2. *Let $W \subset X^2$ be a complete polar cone and (Y, X) be a pre-dual pair for a reflexive semiring $\mathcal{K}$ such that Y separates X . If $(s, t) \notin W$, then, there exists $y \in Y$ such that*

$$(f, g) \in W \implies \langle y \mid f \rangle \leq \langle y \mid g \rangle$$

and

$$\langle y \mid s \rangle \not\leq \langle y \mid t \rangle .$$

Given a pre-dual pair (Y, X) , we define the following correspondences between subsemimodules of X^2 and Y :

$$W \subset X^2 \mapsto W^\circ := \{y \in Y \mid \langle y \mid f \rangle \leq \langle y \mid g \rangle , \forall (f, g) \in W\} ,$$

$$V \subset Y \mapsto V^\circ := \{(f, g) \in X^2 \mid \langle y \mid f \rangle \leq \langle y \mid g \rangle , \forall y \in V\} ,$$

and

$$W \subset X^2 \mapsto W^\top := \{y \in Y \mid \langle y \mid f \rangle = \langle y \mid g \rangle , \forall (f, g) \in W\} ,$$

$$V \subset Y \mapsto V^\perp := \{(f, g) \in X^2 \mid \langle y \mid f \rangle = \langle y \mid g \rangle , \forall y \in V\} .$$

Then, taking $Y = X^{\text{op}}$ and $\langle y \mid x \rangle = x \backslash y$, the universal separation theorem for complete semimodules of Section 2 (see also [CGQ04, Th. 13]) implies that the following statements are equivalent:

- (i) $V \subset Y$ is a complete semimodule,
- (ii) $V = (V^\circ)^\circ$,
- (iii) $V = (V^\perp)^\top$.

Now, thanks to the previous separation theorems, we can prove the dual result.

Theorem 7. *Let (Y, X) be a pre-dual pair which satisfies the following property: If $W \subset X^2$ is a complete polar cone (resp. complete congruence) and $(s, t) \notin W$, there exists $y \in Y$ such that*

$$(f, g) \in W \implies \langle y \mid f \rangle \leq \langle y \mid g \rangle$$

and

$$\langle y \mid s \rangle \not\leq \langle y \mid t \rangle.$$

Then, a subsemimodule $W \subset X^2$ is a complete polar cone (resp. complete congruence) if, and only if,

$$W = (W^\circ)^\circ \text{ (resp. } W = (W^\top)^\perp \text{)}.$$

Proof. We only prove the theorem for polar cones. The case of congruences is similar.

Note that any set of the form V° is a complete polar cone. This follows from the fact that the map $x \mapsto \langle y \mid x \rangle$ is right linear and continuous for all $y \in Y$.

Since the inclusion $W \subset (W^\circ)^\circ$ is straightforward, it suffices to show that $(W^\circ)^\circ \subset W$. Assume that $(s, t) \notin W$. Then, there exists $y \in Y$ such that $\langle y \mid s \rangle \not\leq \langle y \mid t \rangle$ and $\langle y \mid f \rangle \leq \langle y \mid g \rangle$ for all $(f, g) \in W$. Since this means that $y \in W^\circ$ but $\langle y \mid s \rangle \not\leq \langle y \mid t \rangle$, we conclude that $(s, t) \notin (W^\circ)^\circ$. $\square$

Note that, by Theorems 5 and 6, the condition of the previous theorem is satisfied when $Y = X^{\text{op}}$ and $\langle y \mid x \rangle = x \setminus y$, and, by Corollaries 1 and 2, it is also satisfied when (Y, X) is a pre-dual pair for a reflexive semiring $\mathcal{K}$ such that Y separates X .

For any subset U of X^2 , we denote by $\overline{\text{pol}}(U)$ (resp. $\overline{\text{cong}}(U)$) the smallest complete polar cone (resp. complete congruence) containing it. Then, we have the following corollary.

Corollary 3. *Let (Y, X) be a pre-dual pair which satisfies the condition in Theorem 7. Then, for any subset U of X^2 we have $\overline{\text{pol}}(U) = (U^\circ)^\circ$ (resp. $\overline{\text{cong}}(U) = (U^\top)^\perp$).*

Proof. Since any set of the form V° is a complete polar cone, it follows that $(U^\circ)^\circ$ is a complete polar cone which clearly contains U .

Let W be a complete polar cone containing U . Then,

$$U \subset W \implies W^\circ \subset U^\circ \implies (U^\circ)^\circ \subset (W^\circ)^\circ,$$

and by Theorem 7 we have $(W^\circ)^\circ = W$. Therefore, $(U^\circ)^\circ$ is the smallest complete polar cone containing U .

The case of complete congruences can be proved along the same lines. $\square$

Example 4. When $\mathcal{K} = \overline{\mathbb{R}}_{\max}$ and $X = \overline{\mathbb{R}}_{\max}^n$, note that if we consider the bracket $\langle y \mid x \rangle := \oplus_i y_i x_i$, Corollary 2 can be stated as follows: If $W \subset (\overline{\mathbb{R}}_{\max}^n)^2$ is a complete polar cone and $s, t \in \overline{\mathbb{R}}_{\max}^n$ are such that $(s, t) \notin W$, then, there exists $y \in \overline{\mathbb{R}}_{\max}^n$ such that

$$(f, g) \in W \implies \oplus_i y_i f_i \leq \oplus_j y_j g_j$$

and

$$\oplus_i y_i s_i \not\leq \oplus_j y_j t_j.$$

For instance, consider the polar cone $W := V^\circ$, where V is the subsemimodule of $\overline{\mathbb{R}}_{\max}^3$ generated by the vectors $a = (2, 0, 3)^T$, $b = (2, 1, 0)^T$ and $c = (2, 4, 0)^T$. This

- If $f_2 \leq 0$ and $f_3 > 2$, then $H_f = \{x \in \overline{\mathbb{R}}_{\max}^3 \mid f_1 x_1 \oplus f_3 x_3 \leq x_2\}$. Note that $(2, 0, 3)^T \notin H_f$ because $f_3 > 2$. Therefore, like in the previous case, the set H_f never contains V because in particular it does not contain the vector a .
- If $f_2 > 0$ and $f_3 > 2$, the elements of H_f cannot have finite second and third entries. Then, in this case, V is never contained in H_f .

Therefore, we conclude that $\hat{t} = (0, 0, 2)^T$ and so $^{-}\hat{t} = -\hat{t} = (0, 0, -2)^T$. Since $^{-}\hat{t} = (-2)b \oplus (-4)c \in V$, it follows that $\langle ^{-}\hat{t} \mid f \rangle \leq \langle ^{-}\hat{t} \mid g \rangle$ for all $(f, g) \in W$. However, $\langle ^{-}\hat{t} \mid s \rangle = 1 > 0 = \langle ^{-}\hat{t} \mid t \rangle$ and thus $^{-}\hat{t}$ separates (s, t) from W .

A natural way to define a congruence W is to take a continuous right linear map F from X to a complete right $\mathcal{K}$ -semimodule Z and to define $W = \ker F := \{(u, v) \in X^2 \mid F(u) = F(v)\}$. Then, since F is continuous, W is complete and it satisfies

$$\hat{u} = \vee \{v \in X \mid F(v) = F(u)\} = \vee \{v \in X \mid F(v) \leq F(u)\} = F^\sharp(F(u)),$$

for all $u \in X$. Conversely, it can be checked that every complete congruence W arises in this way. It suffices to take for F the canonical morphism from X to its quotient by the equivalence relation W . The details are left to the reader.

Remark 3. A basic special case is when $X = \overline{\mathbb{R}}_{\max}^n$ and F is a continuous (right) linear map from X to $\overline{\mathbb{R}}_{\max}^p$, so that F can be represented by a $p \times n$ matrix $A = (a_{ij})$ with entries in $\overline{\mathbb{R}}_{\max}$, meaning that the i -th coordinate of F is given by $F_i(x) = \max_j (a_{ij} + x_j)$. The residual of F is the min-plus linear map represented by the matrix A transposed and changed of sign $(-a_{ji})$, $F_j^\sharp(y) = \min_i (-a_{ij} + y_i)$, with the convention that $+\infty$ is absorbing for the (usual) addition. Hence, the maximal representative $\hat{u} = F^\sharp \circ F(u)$ is determined by a simple “min-max” combinatorial formula.

4. SEPARATION THEOREMS FOR THE MAX-PLUS SEMIRING

In the preceding section, the semimodules, polar cones, and congruences were required to be complete, which is the same as being closed in the Scott topology. This topology is adapted when the underlying semiring is the completed max-plus semiring, $\overline{\mathbb{R}}_{\max}$ (we refer the reader to [Aki99, AS03] for a discussion of this topology within the max-plus setting). However, in many applications, the semiring of interest is rather the non completed max-plus semiring $\mathbb{R}_{\max}$, and the topology of choice is the standard one, which can be defined by the metric $d(a, b) := |\exp(a) - \exp(b)|$. Thus, instead of complete pre-congruences over $\overline{\mathbb{R}}_{\max}^n$, we now deal with pre-congruences over $\mathbb{R}_{\max}^n$ that are closed (in the induced product topology), and we would like to find separation theorems in the spirit of Theorem 5 involving closed separating sets. We may of course consider the restriction to $(\mathbb{R}_{\max}^n)^2$ of the separating set constructed in the proof of this theorem, i.e.

$$\{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid g \setminus \hat{t} \leq f \setminus \hat{t}\},$$

but it can be checked that this set is not closed as soon as $\hat{t}$ has a $-\infty$ coordinate (see the example below). In this section, we apply a perturbation technique to derive separation theorems which are appropriate for closed polar cones and congruences over $\mathbb{R}_{\max}^n$.

Example 5. Consider again the polar cone $W = V^\circ$ and the pair of vectors (s, t) of Example 4, but assume that we add the vector $(e, \varepsilon, \varepsilon)^T$ to the set of generators of

V. Then, it can be checked that with this modification we have $\hat{t} = t = (\varepsilon, 0, 2)^T$. Therefore, the restriction to $(\mathbb{R}_{\max}^3)^2$ of the set that separates (s, t) from W (given by Theorem 5) is

$$\begin{aligned} & \{(f, g) \in (\mathbb{R}_{\max}^3)^2 \mid (+\infty)f_1 \oplus f_2 \oplus (-2)f_3 \leq (+\infty)g_1 \oplus g_2 \oplus (-2)g_3\} = \\ & \{(f, g) \in (\mathbb{R}_{\max}^3)^2 \mid g_1 \neq \varepsilon\} \cup \\ & \{(f, g) \in (\mathbb{R}_{\max}^3)^2 \mid f_1 = g_1 = \varepsilon, f_2 \oplus (-2)f_3 \leq g_2 \oplus (-2)g_3\} \end{aligned}$$

which is not a closed subset of $(\mathbb{R}_{\max}^3)^2$.

Inspired by the previous section, if $W \subset (\mathbb{R}_{\max}^n)^2$ is a closed pre-congruence, for $g \in \mathbb{R}_{\max}^n$ we set

$$(10) \quad \bar{g} := \sup\{f \in \mathbb{R}_{\max}^n \mid (f, g) \in W\} \in \overline{\mathbb{R}}_{\max}^n.$$

Unlike the case of complete pre-congruences, observe that $\bar{g}$ may have some entries equal to $+\infty$, so $(\bar{g}, g)$ need not belong to W . However, since $(g, g) \in W$, we still have $g \leq \bar{g}$ for all $g \in \mathbb{R}_{\max}^n$. We shall use the fact that

$$(11) \quad g \leq f \implies \bar{g} \leq \bar{f},$$

which can be proved as in the case of complete pre-congruences.

Due to the fact that $(\bar{g}, g)$ does not necessarily belong to W , we cannot use the vector $\bar{g}$ in the same way we did it before. Therefore, we need a perturbation argument in which the following simple construction will be very important. For $z \in \overline{\mathbb{R}}_{\max}^n$ and $\beta \in \mathbb{R}_{\max}$, we define the vector $z^\beta \in \mathbb{R}_{\max}^n$ by:

$$z_i^\beta = \begin{cases} \beta & \text{if } z_i = +\infty, \\ z_i & \text{otherwise.} \end{cases}$$

In other words, z^β is obtained from z by replacing its $+\infty$ entries by β . We denote by e_i the i -th unit vector, i.e. the vector defined by $(e_i)_i := e$ and $(e_i)_j := \varepsilon$ if $j \neq i$. We shall need the following lemma.

Lemma 2. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a congruence and $f, g \in \mathbb{R}_{\max}^n$ be such that $(f, g) \in W$. Then, $(g, g \oplus e_i \gamma) \in W$ for all $\gamma \leq f_i$.*

Proof. If $\gamma \leq f_i$, since W is a congruence and $(f, g) \in W$, we have $(f, g \oplus e_i \gamma) = (f \oplus e_i \gamma, g \oplus e_i \gamma) = (f, g) \oplus (e_i, e_i) \gamma \in W$. Hence, $(g, g \oplus e_i \gamma) \in W$ because $(g, f) \in W$, $(f, g \oplus e_i \gamma) \in W$ and W is a congruence. $\square$

Then we have the following corollary.

Corollary 4. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed congruence or a closed polar cone. Then, for all $g \in \mathbb{R}_{\max}^n$, if $\beta \geq \max_i(g_i)$ we have $(\bar{g}^\beta, g) \in W$.*

Proof. Let us first consider the case where W is a closed congruence. If $\bar{g}_i = +\infty$, there exists $f \in \mathbb{R}_{\max}^n$ such that $(f, g) \in W$ and $f_i > \beta$. Then, by Lemma 2 we have $(g, g \oplus e_i \beta) \in W$.

Now assume that $\bar{g}_i < +\infty$. Then, if $\gamma < \bar{g}_i$, there exists $f \in \mathbb{R}_{\max}^n$ such that $(f, g) \in W$ and $f_i > \gamma$. Thus, by Lemma 2 we have $(g, g \oplus e_i \gamma) \in W$. Since this holds for all $\gamma \in \mathbb{R}_{\max}$ such that $\gamma < \bar{g}_i$, it follows that $(g, g \oplus e_i \bar{g}_i) \in W$ because W is closed.

Consequently, as W is a semimodule, we have

$$\begin{aligned} (\bar{g}^\beta, g) &= ((\oplus_{\{i \mid \bar{g}_i < +\infty\}} g \oplus e_i \bar{g}_i) \oplus (\oplus_{\{i \mid \bar{g}_i = +\infty\}} g \oplus e_i \beta), g) \\ &= (\oplus_{\{i \mid \bar{g}_i < +\infty\}} (g \oplus e_i \bar{g}_i, g)) \oplus (\oplus_{\{i \mid \bar{g}_i = +\infty\}} (g \oplus e_i \beta, g)) \in W. \end{aligned}$$

The case of polar cones is easier (indeed the assertion holds for any $\beta \in \mathbb{R}_{\max}$). If $\bar{g}_i = +\infty$, there exists $f \in \mathbb{R}_{\max}^n$ such that $(f, g) \in W$ and $f_i > \beta$. Then, as $e_i \beta \leq f$ and $(f, g) \in W$, it follows that $(e_i \beta, g) \in W$, because W is a polar cone.

Assume now that $\bar{g}_i < +\infty$. Then, if $\alpha < \bar{g}_i$, there exists $f \in \mathbb{R}_{\max}^n$ such that $(f, g) \in W$ and $f_i > \alpha$. Therefore, as $e_i \alpha \leq f$ and $(f, g) \in W$, we have $(e_i \alpha, g) \in W$. Since this holds for all $\alpha \in \mathbb{R}_{\max}$ such that $\alpha < \bar{g}_i$, we deduce that $(e_i \bar{g}_i, g) \in W$, because W is closed.

Finally, as W is a semimodule, we get

$$\begin{aligned} (\bar{g}^\beta, g) &= ((\oplus_{\{i \mid \bar{g}_i < +\infty\}} e_i \bar{g}_i) \oplus (\oplus_{\{i \mid \bar{g}_i = +\infty\}} e_i \beta), g) \\ &= ((\oplus_{\{i \mid \bar{g}_i < +\infty\}} (e_i \bar{g}_i, g)) \oplus (\oplus_{\{i \mid \bar{g}_i = +\infty\}} (e_i \beta, g))) \in W, \end{aligned}$$

which completes the proof of the corollary. $\square$

Like in the case of complete pre-congruences, we have:

Lemma 3. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed congruence or a closed polar cone. Then,*

$$(f, g) \in W \implies g \setminus \bar{h} \leq f \setminus \bar{h}$$

for all $h \in \mathbb{R}_{\max}^n$.

Proof. When $g \setminus \bar{h} = -\infty$ the assertion is obvious, so assume that $g \setminus \bar{h} \neq -\infty$. Let $\alpha < g \setminus \bar{h}$, so that $\bar{h} \geq g\alpha$. Then, if $\beta \geq \max_i (g_i \alpha \oplus h_i)$, we have $g\alpha \leq \bar{h}^\beta$, and by Corollary 4 we know that $(\bar{h}^\beta, h) \in W$. Since W is a pre-congruence, from $(\bar{h}^\beta \oplus f\alpha, \bar{h}^\beta) = (\bar{h}^\beta \oplus f\alpha, \bar{h}^\beta \oplus g\alpha) = (\bar{h}^\beta, \bar{h}^\beta) \oplus (f, g)\alpha \in W$ and $(\bar{h}^\beta, h) \in W$, it follows that $(\bar{h}^\beta \oplus f\alpha, h) \in W$, and so, $f\alpha \leq \bar{h}^\beta \oplus f\alpha \leq \bar{h}$. Therefore, we have $f \setminus \bar{h} \geq \alpha$. Since this holds for all $\alpha \in \mathbb{R}_{\max}$ such that $\alpha < g \setminus \bar{h}$, we conclude that $g \setminus \bar{h} \leq f \setminus \bar{h}$. $\square$

The following lemmas will be useful to prove the separation theorems for the max-plus semiring.

Lemma 4. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed pre-congruence. Assume that $s, t, h \in \mathbb{R}_{\max}^n$ are such that $t \leq h$ and $s \not\leq \bar{h}$. Then, we have $t \setminus \bar{h} \not\leq s \setminus \bar{h}$.*

Proof. Assume that $t \setminus \bar{h} \leq s \setminus \bar{h}$. Then, we get $s(t \setminus \bar{h}) \leq s(s \setminus \bar{h}) \leq \bar{h}$. Since $t \leq h$, by (11) we have $t \leq \bar{t} \leq \bar{h}$, and so $e \leq t \setminus \bar{h}$. Therefore, it follows that $s = se \leq s(t \setminus \bar{h}) \leq \bar{h}$, which is a contradiction. $\square$

Lemma 5. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed congruence or a closed polar cone. Assume that $s, t \in \mathbb{R}_{\max}^n$ are such that $s \not\leq \bar{t}$. Then, there exists $h \in \mathbb{R}_{\max}^n$ with finite entries such that $t \leq h$ and $s \not\leq \bar{h}$.*

Proof. Assume that $s \leq \bar{h}$ for all $h \in \mathbb{R}_{\max}^n$ with finite entries that satisfy $t \leq h$. Let $\{h_k\}_{k \in \mathbb{N}} \subset \mathbb{R}_{\max}^n$ be a decreasing sequence of vectors with finite entries such that $\lim_{k \rightarrow \infty} h_k = t$. Then, as by (11) the sequence $\{\bar{h}_k\}_{k \in \mathbb{N}} \subset \overline{\mathbb{R}_{\max}^n}$ is also decreasing, there exists $z \in \overline{\mathbb{R}_{\max}^n}$ such that $\lim_{k \rightarrow \infty} \bar{h}_k = z$. Since $s \leq \bar{h}_k$ for all $k \in \mathbb{N}$, we have $s \leq z$. Note that we can assume, without loss of generality, that $(\bar{h}_k)_i = +\infty \iff z_i = +\infty$. If $\lambda \geq \max_i (h_1)_i$, then by Corollary 4 we know that $((\bar{h}_k)^\lambda, h_k) \in W$ for all $k \in \mathbb{N}$. Therefore, we get $(z^\lambda, t) = \lim_{k \rightarrow \infty} ((\bar{h}_k)^\lambda, h_k) \in W$,

because W is closed. Since this holds for all $\lambda \in \mathbb{R}_{\max}$ such that $\lambda \geq \max_i (h_1)_i$, it follows that $\bar{t} \geq z \geq s$, which contradicts our assumption. $\square$

As a consequence of the previous results we obtain the separation theorem for closed polar cones.

Theorem 8. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed polar cone. Assume that $s, t \in \mathbb{R}_{\max}^n$ are such that $(s, t) \notin W$. Then, there exists $y \in \mathbb{R}_{\max}^n$ such that*

$$(f, g) \in W \implies \oplus_i f_i y_i \leq \oplus_j g_j y_j$$

and

$$\oplus_i s_i y_i \not\leq \oplus_j t_j y_j .$$

Proof. In the first place, we claim that $s \not\leq \bar{t}$. Indeed, if $s \leq \bar{t}$, let $\beta \geq \max_i (s_i \oplus t_i)$. Then, $s \leq \bar{t}^\beta$ and by Corollary 4 we have $(\bar{t}^\beta, t) \in W$. It follows that $(s, t) \in W$ because W is a polar cone, which contradicts our assumption. This proves our claim.

Since $s \not\leq \bar{t}$, according to Lemma 5 there exists $h \in \mathbb{R}_{\max}^n$ with finite entries such that $t \leq h$ and $s \not\leq \bar{h}$. Therefore, from Lemmas 3 and 4, it follows that

$$(f, g) \in W \implies g \setminus \bar{h} \leq f \setminus \bar{h}$$

and

$$t \setminus \bar{h} \not\leq s \setminus \bar{h} .$$

Then, as $u \setminus \bar{h} = -(\oplus_i u_i (-\bar{h}_i))$ and $\bar{h}_i \geq h_i > -\infty$ for all i , the assertion of the theorem holds with $y := -\bar{h}$. $\square$

In order to prove a separation theorem for closed congruences, we need the following result.

Lemma 6. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed congruence. Then, for all $f, g \in \mathbb{R}_{\max}^n$, the following properties are satisfied.*

$$(12) \quad (f, g) \in W \iff \bar{f} = \bar{g} ;$$

$$(13) \quad f \leq \bar{g} \implies \bar{f} \leq \bar{g} .$$

Proof. (12) If $(f, g) \in W$, we have

$$\{h \in \mathbb{R}_{\max}^n \mid (h, f) \in W\} = \{h \in \mathbb{R}_{\max}^n \mid (h, g) \in W\}$$

because W is a congruence. Therefore, $\bar{f} = \bar{g}$.

Conversely, assume that $\bar{f} = \bar{g}$. If $\lambda \geq \max_i (f_i \oplus g_i)$, by Corollary 4 it follows that $(\bar{f}^\lambda, f) \in W$ and $(\bar{g}^\lambda, g) \in W$. Since $\bar{f} = \bar{g}$, we have $\bar{f}^\lambda = \bar{g}^\lambda$, and thus $(f, g) \in W$ because W is a congruence.

(13) Let $\lambda \geq \max_i (f_i \oplus g_i)$. Then, if $f \leq \bar{g}$ we have $f \leq \bar{g}^\lambda$, and by Corollary 4 we know that $(\bar{g}^\lambda, g) \in W$. Therefore, by (11) and (12), it follows that $\bar{f} \leq \bar{g}^\lambda = \bar{g}$. $\square$

Theorem 9. *Let $W \subset (\mathbb{R}_{\max}^n)^2$ be a closed congruence. Assume that $s, t \in \mathbb{R}_{\max}^n$ are such that $(s, t) \notin W$. Then, there exists $y \in \mathbb{R}_{\max}^n$ such that*

$$(f, g) \in W \implies \oplus_i f_i y_i = \oplus_j g_j y_j$$

and

$$\oplus_i s_i y_i \neq \oplus_j t_j y_j .$$

Proof. Since $(s, t) \notin W$, by (12) we have $\bar{s} \neq \bar{t}$, hence, $\bar{s} \not\leq \bar{t}$ and/or $\bar{t} \not\leq \bar{s}$. Assume for instance that $\bar{s} \not\leq \bar{t}$. Then, from (13) it follows that $s \not\leq \bar{t}$. Therefore, according to Lemma 5 there exists $h \in \mathbb{R}_{\max}^n$ with finite entries such that $t \leq h$ and $s \not\leq \bar{h}$. Since $(f, g) \in W$ implies $(g, f) \in W$ because W is a congruence, from Lemmas 3 and 4, we have

$$(f, g) \in W \implies f \setminus \bar{h} = g \setminus \bar{h}$$

and

$$t \setminus \bar{h} \neq s \setminus \bar{h}.$$

Then, taking into account that $u \setminus \bar{h} = -(\oplus_i u_i(-\bar{h}_i))$ and $\bar{h}_i \geq h_i > -\infty$ for all i , the assertion of the theorem holds with $y := -\bar{h}$. $\square$

Like in the previous section, consider the following correspondences between subsemimodules of $(\mathbb{R}_{\max}^n)^2$ and $\mathbb{R}_{\max}^n$:

$$W \subset (\mathbb{R}_{\max}^n)^2 \mapsto W^\circ := \{x \in \mathbb{R}_{\max}^n \mid \oplus_i f_i x_i \leq \oplus_j g_j x_j, \forall (f, g) \in W\},$$

$$V \subset \mathbb{R}_{\max}^n \mapsto V^\circ := \{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid \oplus_i f_i x_i \leq \oplus_j g_j x_j, \forall x \in V\},$$

and

$$W \subset (\mathbb{R}_{\max}^n)^2 \mapsto W^\top := \{x \in \mathbb{R}_{\max}^n \mid \oplus_i f_i x_i = \oplus_j g_j x_j, \forall (f, g) \in W\},$$

$$V \subset \mathbb{R}_{\max}^n \mapsto V^\perp := \{(f, g) \in (\mathbb{R}_{\max}^n)^2 \mid \oplus_i f_i x_i = \oplus_j g_j x_j, \forall x \in V\}.$$

Then, by the separation theorem for closed semimodules (see [Zim77, Th. 4], [SS92], see also [CGQS05, Th. 3.14] for recent improvements), it follows that the following statements are equivalent:

- (i) $V \subset \mathbb{R}_{\max}^n$ is a closed semimodule,
- (ii) $V = (V^\circ)^\circ$,
- (iii) $V = (V^\perp)^\top$.

Now, as a consequence of Theorems 8 and 9, we obtain the following dual result. We omit the proof because it is similar to that of Theorem 7.

Theorem 10. *A semimodule $W \subset (\mathbb{R}_{\max}^n)^2$ is a closed polar cone (resp. closed congruence) if, and only if,*


$$W = (W^\circ)^\circ \text{ (resp. } W = (W^\top)^\perp \text{)}.$$

As a corollary of the previous theorem, it follows that for any subset U of $(\mathbb{R}_{\max}^n)^2$, $(U^\circ)^\circ$ (resp. $(U^\top)^\perp$) is the smallest closed polar cone (resp. closed congruence) containing it.

Example 6. Consider the semimodule $V \subset \mathbb{R}_{\max}^3$ generated by the vectors $a = (0, 0, 0)^T$, $b = (0, 1, -1)^T$ and $c = (0, 2, -2)^T$. This semimodule is represented by the bounded dark gray region of Figure 3 together with the segments joining the points a and c to it. Then V° is the solution set of a system of homogeneous max-plus linear inequalities, and hence also of a system of homogeneous max-plus linear equations. More precisely,

$$\begin{aligned} V^\circ &= \{(f, g) \in (\mathbb{R}_{\max}^3)^2 \mid \oplus_i f_i x_i \leq \oplus_j g_j x_j, x = a, b, c\} \\ &= \{(f, g) \in (\mathbb{R}_{\max}^3)^2 \mid \oplus_i (f_i \oplus g_i) x_i = \oplus_j g_j x_j, x = a, b, c\}. \end{aligned}$$

We know that it is a finitely generated subsemimodule of $\mathbb{R}_{\max}^6$ (see [BH84, Gau92, GP97]). Solving this system by the elimination method (see [BH84] and [AGG08])

FIGURE 3. Valid linear inequality and equality for V .

for recent improvements), we obtain that the polar cone V° is the subsemimodule of $\mathbb{R}_{\max}^6$ generated by the columns of the following matrix

$$\begin{pmatrix} \varepsilon & \varepsilon & \varepsilon & 1 & \varepsilon & 2 & 0 & 0 & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon \\ 0 & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon \\ \varepsilon & 0 & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon \\ 2 & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon \\ \varepsilon & 0 & \varepsilon & 0 & \varepsilon & 0 & 0 & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & 0 & \varepsilon \\ \varepsilon & \varepsilon & \varepsilon & 1 & 4 & 3 & \varepsilon & 2 & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & 0 \end{pmatrix}.$$

The ordering of the variables in the system of equations above has been chosen such that, if we denote by f the vector whose entries are the first three entries of a solution and by g the vector whose entries are the remaining three entries, then $(f, g) \in V^\circ$. For instance, the fourth column of the matrix above corresponds to the pair of vectors $f = (1, \varepsilon, \varepsilon)^T$ and $g = (\varepsilon, 0, 1)^T$. With each pair of vectors $(f, g) \in V^\circ$ is associated the linear inequality $\oplus_i f_i x_i \leq \oplus_j g_j x_j$ which is said to be *valid* for V because it is satisfied by all its elements. For the previous pair of vectors, we have the valid inequality $1x_1 \leq x_2 \oplus 1x_3$, which is represented by the unbounded light gray region on the left-hand side of Figure 3. Analogously, the congruence $V^\perp$ is the subsemimodule of $\mathbb{R}_{\max}^6$ generated by the columns of the following matrix

$$\begin{pmatrix} 0 & 1 & \varepsilon & 0 & 2 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & 2 & 2 & 0 & 0 & 0 & \varepsilon & \varepsilon \\ 0 & 0 & 0 & \varepsilon & \varepsilon & 0 & 0 & 0 & 0 & \varepsilon & 0 & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon \\ \varepsilon & \varepsilon & 0 & 0 & \varepsilon & 4 & 3 & \varepsilon & \varepsilon & 2 & 1 & 4 & \varepsilon & 3 & \varepsilon & 2 & \varepsilon & \varepsilon & 0 \\ \varepsilon & \varepsilon & \varepsilon & 0 & 2 & \varepsilon & 2 & 0 & \varepsilon & 0 & 1 & \varepsilon & 2 & \varepsilon & 0 & \varepsilon & 0 & \varepsilon & \varepsilon \\ 0 & 0 & 0 & \varepsilon & 0 & \varepsilon & \varepsilon & 0 & 0 & \varepsilon & 0 & 0 & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon \\ \varepsilon & 1 & \varepsilon & \varepsilon & \varepsilon & 4 & 3 & \varepsilon & 0 & 2 & \varepsilon & 4 & \varepsilon & 3 & 0 & 2 & \varepsilon & \varepsilon & 0 \end{pmatrix}.$$

In this case, to solve the system of equations for $V^\perp$, we have taken the same ordering of the variables as for V° . Then, for example, the first column corresponds to the equality $x_1 \oplus x_2 = x_2$ which is equivalent to the valid inequality $x_1 \leq x_2$ (the seventh column of the matrix containing the generators of V°).

Note that, as mentioned in the introduction, any valid inequality $\oplus_i f_i x_i \leq \oplus_j g_j x_j$ for V implies a valid equality for V (meaning that it is satisfied by all its elements), namely $\oplus_i (f_i \oplus g_i) x_i = \oplus_j g_j x_j$. However, not all the valid equalities

for V can be obtained in this way. For instance, the equality $x_2 \oplus 3x_3 = 2x_1 \oplus 3x_3$ is satisfied by all the elements of V . It corresponds to the seventh column of the latter matrix and it is represented by the unbounded light gray region on the right-hand side of Figure 3. This equality cannot be obtained as a max-plus linear combination of the equalities that can be derived from the inequalities in V° .

Acknowledgment. The authors would like to thank the anonymous referees for their valuable comments and suggestions, which improved this manuscript.

REFERENCES

- [AGG08] X. Allamigeon, S. Gaubert, and É. Goubault. Inferring min and max invariants using max-plus polyhedra. In *Proceedings of the 15th International Static Analysis Symposium (SAS'08)*, volume 5079 of *Lecture Notes in Comput. Sci.*, pages 189–204. Springer, 2008.
- [Aki99] M. Akian. Densities of idempotent measures and large deviations. *Trans. Amer. Math. Soc.*, 351(11):4515–4543, 1999.
- [AR06] G. Adilov and A. Rubinov. $\mathbb{B}$ -convex sets and functions. *Numer. Funct. Anal. Optim.*, 27:237–257, 2006.
- [AS03] M. Akian and I. Singer. Topologies on lattice ordered groups, separation from closed downward sets and conjugations of type Lau. *Optimization*, 52(6):629–672, 2003.
- [BCOQ92] F. Baccelli, G. Cohen, G. J. Olsder, and J. P. Quadrat. *Synchronization and Linearity*. Wiley, Chichester, 1992.
- [BH84] P. Butkovič and G. Hegedüs. An elimination method for finding all solutions of the system of linear equations over an extremal algebra. *Ekonomicko-matematicky Obzor*, 20(2):203–215, 1984.
- [BH04] W. Briec and C. Horvath. $\mathbb{B}$ -convexity. *Optimization*, 53:103–127, 2004.
- [BHR05] W. Briec, C. Horvath, and A. Rubinov. Separation in $\mathbb{B}$ -convexity. *Pacific Journal of Optimization*, 1:13–30, 2005.
- [BJ72] T. S. Blyth and M. F. Janowitz. *Residuation Theory*. Pergamon Press, London, 1972.
- [Bou64] N. Bourbaki. *Espaces Vectoriels Topologiques*. Éléments de Mathématique, Livre V. Hermann, 1964.
- [BSS07] P. Butkovič, H. Schneider, and S. Sergeev. Generators, extremals and bases of max cones. *Linear Algebra and Appl.*, 421:394–406, 2007. Also e-print arXiv:math.RA/0604454.
- [CDQV85] G. Cohen, D. Dubois, J. P. Quadrat, and M. Viot. A linear system theoretic view of discrete event processes and its use for performance evaluation in manufacturing. *IEEE Trans. on Automatic Control*, AC-30:210–220, 1985.
- [CGQ99] G. Cohen, S. Gaubert, and J. P. Quadrat. Max-plus algebra and system theory: where we are and where to go now. *Annual Reviews in Control*, 23:207–219, 1999.
- [CGQ01] G. Cohen, S. Gaubert, and J. P. Quadrat. Hahn-Banach separation theorem for max-plus semimodules. In J. L. Menaldi, E. Rofman, and A. Sulem, editors, *Optimal Control and Partial Differential Equations*, pages 325–334. IOS Press, 2001.
- [CGQ04] G. Cohen, S. Gaubert, and J. P. Quadrat. Duality and separation theorems in idempotent semimodules. *Linear Algebra and Appl.*, 379:395–422, 2004. Also e-print arXiv:math.FA/0212294.
- [CGQS05] G. Cohen, S. Gaubert, J. P. Quadrat, and I. Singer. Max-plus convex sets and functions. In G. L. Litvinov and V. P. Maslov, editors, *Idempotent Mathematics and Mathematical Physics*, Contemporary Mathematics, pages 105–129. American Mathematical Society, 2005. Also ESI Preprint 1341, arXiv:math.FA/0308166.
- [DS04] M. Develin and B. Sturmfels. Tropical convexity. *Doc. Math.*, 9:1–27 (electronic), 2004. Also e-print arXiv:math.MG/0308254.
- [DY07] M. Develin and J. Yu. Tropical polytopes and cellular resolutions. *Experimental Mathematics*, 16(3):277–292, 2007. Also e-print arXiv:math.CO/0605494.
- [Gau92] S. Gaubert. *Théorie des systèmes linéaires dans les dioïdes*. Thèse, École des Mines de Paris, July 1992.

- [GHK⁺03] G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. Mislove, and D. S. Scott. *Continuous lattices and domains*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2003.
- [GK04] S. Gaubert and R. D. Katz. Rational semimodules over the max-plus semiring and geometric approach to discrete event systems. *Kybernetika*, 40(2):153–180, 2004. Also e-print arXiv:math.OC/0208014.
- [GK06] S. Gaubert and R. D. Katz. Max-plus convex geometry. In R. A. Schmidt, editor, *Proceedings of the 9th International Conference on Relational Methods in Computer Science and 4th International Workshop on Applications of Kleene Algebra (RelMiCS/AKA 2006)*, volume 4136 of *Lecture Notes in Comput. Sci.*, pages 192–206. Springer, 2006.
- [GK07] S. Gaubert and R. D. Katz. The Minkowski theorem for max-plus convex sets. *Linear Algebra and Appl.*, 421:356–369, 2007. Also e-print arXiv:math.GM/0605078.
- [GM08] S. Gaubert and F. Meunier. Carathéodory, Helly and the others in the max-plus world. *Discrete Computational Geometry*, 2009. In press (published on line), also e-print arXiv:0804.1361.
- [GP97] S. Gaubert and M. Plus. Methods and applications of $(\max, +)$ linear algebra. In R. Reischuk and M. Morvan, editors, *14th Symposium on Theoretical Aspects of Computer Science, STACS 97 (Lübeck)*, volume 1200 of *Lecture Notes in Comput. Sci.*, pages 261–282. Springer, Berlin, 1997.
- [GS07] S. Gaubert and S. Sergeev. Cyclic projectors and separation theorems in idempotent convex geometry. *Journal of Mathematical Sciences*, 155(6):815–829, 2008. Russian version published in *Fundamentalnaya i prikladnaya matematika*, 13(4):33–52, 2007. Also e-print arXiv:0706.3347.
- [Hel88] S. Helbig. On Caratheodory’s and Kreĭn-Milman’s theorems in fully ordered groups. *Comment. Math. Univ. Carolin.*, 29(1):157–167, 1988.
- [Jos05] M. Joswig. Tropical halfspaces. In *Combinatorial and computational geometry*, volume 52 of *Math. Sci. Res. Inst. Publ.*, pages 409–431. Cambridge Univ. Press, Cambridge, 2005. Also e-print arXiv:math.CO/0312068.
- [JSY07] M. Joswig, B. Sturmfels, and J. Yu. Affine buildings and tropical convexity. *Albanian Journal of Mathematics*, 1(4):187–211, 2007. Also e-print arXiv:0706.1918.
- [Kat07] R. D. Katz. Max-plus (A, B) -invariant spaces and control of timed discrete event systems. *IEEE Trans. on Automatic Control*, 52(2):229–241, 2007. Also e-print arXiv:math.OC/0503448.
- [KM97] V. N. Kolokoltsov and V. P. Maslov. *Idempotent analysis and applications*. Kluwer Acad. Publisher, 1997.
- [LGKL08] M. Di Loreto, S. Gaubert, R. D. Katz, and J. J. Loiseau. Duality between invariant spaces for max-plus linear discrete event systems. E-print arXiv:0901.2915, 2008.
- [LMS01] G. L. Litvinov, V. P. Maslov, and G. B. Shpiz. Idempotent functional analysis: An algebraic approach. *Math. Notes*, 69(5):696–729, 2001. Also e-print arXiv:math.FA/0009128.
- [LS07] G. L. Litvinov and G. Shpiz. Kernel theorems and nuclearity in idempotent mathematics. an algebraic approach. *J. of Math. Sci.*, 141(4):1417–1428, 2007. Also e-print arXiv:math.FA/0609033.
- [Mas87] V. P. Maslov. *Méthodes Operatorielles*. Mir, Moscou, 1987.
- [MS92] V. P. Maslov and S. Samborskiĭ, editors. *Idempotent analysis*, volume 13 of *Adv. in Sov. Math.* AMS, RI, 1992.
- [NS07a] V. Nitica and I. Singer. Max-plus convex sets and max-plus semispaces. I. *Optimization*, 56(1–2):171–205, 2007.
- [NS07b] V. Nitica and I. Singer. Max-plus convex sets and max-plus semispaces. II. *Optimization*, 56(3):293–303, 2007.
- [NS07c] V. Nitica and I. Singer. The structure of max-plus hyperplanes. *Linear Algebra and Appl.*, 426:382–414, 2007.
- [Sin97] I. Singer. *Abstract convex analysis*. Wiley, 1997.
- [SS92] S. N. Samborskiĭ and G. B. Shpiz. Convex sets in the semimodule of bounded functions. In *Idempotent analysis*, pages 135–137. Amer. Math. Soc., Providence, RI, 1992.
- [Zim77] K. Zimmermann. A general separation theorem in extremal algebras. *Ekonomicko-matematicheskii Obzor*, 13(2):179–201, 1977.

INRIA AND CENTRE DE MATHÉMATIQUES APPLIQUÉES, ÉCOLE POLYTECHNIQUE. POSTAL ADDRESS: CMAP, ÉCOLE POLYTECHNIQUE, 91128 PALAISEAU CÉDEX, FRANCE. TÈL: +33 1 69 33 46 13, FAX: +33 1 39 63 57 86

E-mail address: `Stephane.Gaubert@inria.fr`

CONICET. POSTAL ADDRESS: INSTITUTO DE MATEMÁTICA “BEPPPO LEVI”, UNIVERSIDAD NACIONAL DE ROSARIO, AVENIDA PELLEGRINI 250, 2000 ROSARIO, ARGENTINA.

E-mail address: `rkatz@fceia.unr.edu.ar`